European Journal of Pharmacology – Molecular Pharmacology Section, 188 (1990) 423–426
Elsevier

Author index

Andersen, P.H. and Jansen, J.A.
Dopamine receptor agonists: selectivity and dopamine D_1 receptor efficacy (188) 335
Arrang, J.-M., Roy, J., Morgat, J.-L., Schunack, W. and Schwartz, J.-C.
Histamine H_3 receptor binding sites in rat brain membranes: modulations by guanine nucleotides and divalent cations (188) 219

Baba, A., see Yara, H. (188) 359
Barefoot, D.S., see Gordon, J.C. (188) 313
Barnes, M.R., see Cowlen, M.S. (188) 105
Bdolah, A., see Galron, R. (188) 85
Belelli, D., see Lan, N.C. (188) 403
Benedetto, P., see Lew, J.Y. (188) 329
Bitran, J.A., Potter, W.Z., Manji, H.K. and Gusovsky, F.
Chronic Li^+ attenuates agonist- and phorbol ester-mediated Na^+/H^+ antiporter activity in HL-60 cells (188) 193
Björklund, P.-E., see Tiger, G. (188) 51
Bockaert, J., see Pin, J.-P. (188) 81
Borden, L.A. and Gibbs, T.T.
Flunitrazepam photoaffinity labeling of the $GABA_A$ receptor reduces inhibition of [^{3}H]Ro15-4513 binding by GABA (188) 391
Boulay, G., see Guillemette, G. (188) 251
Boyd, N., see Luber-Narod, J. (188) 185
Buckley, A.R., see Russell, D.H. (188) 139

Carr, C., Loney, C., Unson, C., Knowler, J. and Milligan, G.
Chronic exposure of rat glioma C6 cells to cholera toxin induces loss of the α-subunit of the stimulatory guanine nucleotide-binding protein (G_s) (188) 203
Cattaneo, E. and Maggi, A.
c-fos induction by estrogen in specific rat brain areas (188) 153
Chen, J.-S., see Lan, N.C. (188) 403
Ciardo, A. and Meldolesi, J.
Multiple actions of SC 38249: the blocker of both voltage-operated and second messenger-operated Ca^{2+} channels also inhibits Ca^{2+} extrusion (188) 417
Compton, R.P., Hood, W.F. and Monahan, J.B.
Evidence for a functional coupling of the NMDA and glycine recognition sites in synaptic plasma membranes (188) 63
Compton, R.P., see Hood, W.F. (188) 43
Cowburn, R.F., see Tiger, G. (188) 51
Cowlen, M.S., Barnes, M.R. and Toews, M.L.
Regulation of histamine H_1 receptor-mediated phosphoinositide hydrolysis by histamine and phorbol esters in DDT_1 MF-2 cells (188) 105
Crowe, P.D., see Russell, D.H. (188) 139

Dagerlind, Å., see Schalling, M. (188) 277
Dauzats, M., see Langin, D. (188) 261
De Chaffoy de Courcelles, D. and De Clerck, F.
Effect of thromboxane A_2 synthetase inhibition, singly and combined with thromboxane A_2/prostaglandin endoperoxide receptor antagonism, on inositol phospholipid turnover and on 5-HT release by washed human platelets (188) 161
De Clerck, F., see De Chaffoy de Courcelles, D. (188) 161
De Graan, P.N.E., see Dekker, L.V. (188) 113
Dekker, L.V., De Graan, P.N.E., Spierenburg, H., De Wit, M., Versteeg, D.H.G. and Gispen, W.H.
Evidence for a relationship between B-50 (GAP-43) and [^{3}H]noradrenaline release in rat brain synaptosomes (188) 113
De Wit, M., see Dekker, L.V. (188) 113
Doi, M., see Yara, H. (188) 359

Ehrlich, M., see Schalling, M. (188) 277

Farese, R.V., see Russell, D.H. (188) 139
Favreau, I., see Guillemette, G. (188) 251
Feigenblum, D., see Lew, J.Y. (188) 329
Ferkany, J.W., see Kaplita, P.V. (188) 175
Filer, D., see Lew, J.Y. (188) 329
Fleming, N., see Laniyonu, A. (188) 171
Ford, W., see Hoyer, D. (188) 9
Fowler, C.J., see Tiger, G. (188) 51
Fraeyman, N., see Vanscheeuwijck, P. (188) 129
Fujii, Y., see Yamashita, A. (188) 229
Fukuo, K., see Morimoto, S. (188) 369

Galron, R., Kloog, Y., Bdolah, A. and Sokolovsky, M.
Different pathways of endothelin/sarafotoxin-stimulated phosphoinositide hydrolysis in myocytes (188) 85
Galzi, J.-L., Mejean, A., Goeldner, M., Hirth, C. and Ilien, B.
Photoaffinity labelling of the μ-opioid receptor is dependent on the nature of the photosensitive group of carfentanil derivatives (188) 321
Garlind, A., see Tiger, G. (188) 51
Gee, K.W., see Lan, N.C. (188) 403
Gibbs, T.T., see Borden, L.A. (188) 391
Gispen, W.H., see Dekker, L.V. (188) 113
Godfrey, P.P., see Watson, S.P. (188) 273

0922-4106/90/$03.50 © 1990 Elsevier Science Publishers B.V. (Biomedical Division)

Goeldner, M., see Galzi, J.-L. (188) 321
Gokita, T., see Karibe, H. (188) 407
Goldstein, M., see Harada, K. (188) 123
Goldstein, M., see Lew, J.Y. (188) 329
Goldstein, M., see Schalling, M. (188) 277
Gordon, J.C., Sarbin, N.S., Barefoot, D.S. and Pinkus, L.M.
Solubilization of a 5-HT_3 binding site from rabbit small bowel muscularis membranes (188) 313
Greengard, P., see Schalling, M. (188) 277
Guillemette, G., Favreau, I., Lamontagne, S. and Boulay, G.
2,3-Diphosphoglycerate is a nonselective inhibitor of inositol 1,4,5-trisphosphate action and metabolism (188) 251
Gusovsky, F., see Bitran, J.A. (188) 193

Hadden, E.M., see Russell, D.H. (188) 139
Harada, K., Meller, E. and Goldstein, M.
Effects of pertussis toxin on inhibition of synaptosomal tyrosine hydroxylase activity by apomorphine (188) 123
Hirth, C., see Galzi, J.-L. (188) 321
Hökfelt, T., see Schalling, M. (188) 277
Hollenberg, M.D., see Yang, S.-G. (188) 289
Hood, W.F., Thomas, J.W., Compton, R.P. and Monahan, J.B.
Guanine nucleotide modulation of [^{3}H]TCP binding to the NMDA receptor complex (188) 43
Hood, W.F., see Compton, R.P. (188) 63
Hoyer, D., Jones, C.R., Ford, W. and Palacios, J.M.
Subtypes of α_1-adrenoceptors in hippocampus of pigs, guinea-pigs, calves and humans: regional differences (188) 9

Ilien, B., see Galzi, J.-L. (188) 321
Imanaka, S., see Morimoto, S. (188) 369
Inoue, M., see Yara, H. (188) 359
Ishibashi, S., see Yamashita, A. (188) 229
Ishida, T., see Yara, H. (188) 359
Iwata, H., see Yara, H. (188) 359

Jansen, J.A., see Andersen, P.H. (188) 335
Johansen, O., see Sager, G. (188) 243
Jones, C.R., see Hoyer, D. (188) 9

Kaplita, P.V. and Ferkany, J.W.
Evidence for direct interactions between the NMDA and glycine recognition sites in brain (188) 175
Karibe, H., Matsuo, K., Gokita, T. and Uchida, M.-K.
Inhibitory effects of protein kinase inhibitors and cytoskeletal inhibitors on Ca^{2+}-free contraction of rat uterus (188) 407
Katagawa, Y., see Yara, H. (188) 359
Kim, S., see Morimoto, S. (188) 369
Kitamura, K., see Yara, H. (188) 359
Kitano, S., see Morimoto, S. (188) 369
Kloog, Y., see Galron, R. (188) 85
Kloog, Y., see Nadler, V. (188) 97
Knowler, J., see Carr, C. (188) 203
Koh, E., see Morimoto, S. (188) 369
Kurokawa, T., see Yamashita, A. (188) 229

Lafontan, M., see Langin, D. (188) 261
Laird II, H.E., see Russell, D.H. (188) 139
Lamontagne, S., see Guillemette, G. (188) 251
Lan, N.C., Chen, J.-S., Belelli, D., Pritchett, D.B., Seeburg, P.H. and Gee, K.W.
A steroid recognition site is functionally coupled to an expressed $GABA_A$-benzodiazepine receptor (188) 403
Langin, D., Paris, H., Dauzats, M. and Lafontan, M.
Discrimination between α_2-adrenoceptors and [^{3}H]idazoxan-labelled non-adrenergic sites in rabbit white fat cells (188) 261
Laniyonu, A., Sliwinski-Lis, E. and Fleming, N.
Muscarinic M_3 receptors are coupled to two signal transduction pathways in rat submandibular cells (188) 171
Leeman, S.E., see Luber-Narod, J. (188) 185
Leissner, J., see Sandmann, J. (188) 89
Lew, J.Y., Zawadzka, H., Feigenblum, D., Tang, D., Filer, D., Benedetto, P. and Goldstein, M.
Regulation and molecular characterization of dopamine D_2 receptors in a prolactin-secreting 7315a anterior pituitary tumor (188) 329
Lindmar, R., see Sandmann, J. (188) 89
Löffelholz, K., see Sandmann, J. (188) 89
Loney, C., see Carr, C. (188) 203
Luber-Narod, J., Boyd, N. and Leeman, S.E.
Guanine nucleotides decrease the affinity of substance P binding to its receptor (188) 185

Maggi, A., see Cattaneo, E. (188) 153
Magnan, J., see Tiberi, M. (188) 379
Manji, H.K., see Bitran, J.A. (188) 193
Marvizón, J.-C.G. and Skolnick, P.
An endogenous modulator of N-methyl-D-aspartate receptor-coupled glycine receptors (188) 23
Matsuo, K., see Karibe, H. (188) 407
McCann, D.J. and Su, T.-P.
Haloperidol-sensitive (+)[^{3}H]SKF-10,047 binding sites (σ sites) exhibit a unique distribution in rat brain subcellular fractions (188) 211
Mejean, A., see Galzi, J.-L. (188) 321
Meldolesi, J., see Ciardo, A. (188) 417
Meller, E., see Harada, K. (188) 123
Milligan, G., see Carr, C. (188) 203
Miyashita, Y., see Morimoto, S. (188) 369
Mjøs, O.D., see Sager, G. (188) 243
Monahan, J.B., see Compton, R.P. (188) 63
Monahan, J.B., see Hood, W.F. (188) 43
Morgat, J.-L., see Arrang, J.-M. (188) 219
Morimoto, S., Kim, S., Fukuo, K., Koh, E., Morita, R., Kitano, S., Miyashita, Y., Imanaka, S. and Ogihara, T.
Participation of both intracellular free Ca^{2+} and protein kinase C in tonic vasoconstriction induced by prostaglandin $F_{2\alpha}$ (188) 369
Morita, R., see Morimoto, S. (188) 369
Murphy, S. and Welk, G.
Hydrolysis of polyphosphoinositides in astrocytes by platelet-activating factor (188) 399

Nadler, V., Kloog, Y. and Sokolovsky, M.
Distinctive structural requirement for the binding of uncompetitive blockers (phencyclidine-like drugs) to the NMDA receptor (188) 97
Nordstedt, C.
Synergism between phorbol ester and the Ca^{2+} ionophore A23187 on protein kinase C translocation, [^{3}H]PDBu binding and adenosine A_2-receptor activation in Jurkat cells (188) 349

Ogihara, T., see Morimoto, S. (188) 369
O'Neill, C., see Tiger, G. (188) 51

Palacios, J.M., see Hoyer, D. (188) 9
Paris, H., see Langin, D. (188) 261
Pettibone, D.J., Woyden, C.J. and Totaro, J.A.
Identification of functional oxytocin receptors in lactating rat mammary gland in vitro (188) 235
Pin, J.-P. and Bockaert, J.
ω-Conotoxin GVIA and dihydropyridines discriminate two types of Ca^{2+} channels involved in GABA release from striatal neurons in culture (188) 81
Pinkus, L.M., see Gordon, J.C. (188) 313
Potter, W.Z., see Bitran, J.A. (188) 193
Pritchett, D.B., see Lan, N.C. (188) 403

Randall, P.K., see Severson, J.A. (188) 283
Reith, M.E.A.
[^{14}C]Guanidinium ion influx into Na^+ channel preparations from mouse cerebral cortex (188) 33
Richardson, J.S., see Tiong, A.H.K. (188) 411
Roy, J., see Arrang, J.-M. (188) 219
Russell, D.H., Zorn, N.E., Buckley, A.R., Crowe, P.D., Sauro, M.D., Hadden, E.M., Farese, R.V. and Laird II, H.E.
Prolactin and known modulators of rat splenocytes activate nuclear protein kinase C (188) 139

Sadatome, T., see Yara, H. (188) 359
Sager, G., Johansen, O. and Mjøs, O.D.
Effects of insulin on mononuclear leukocyte β-adrenoceptor density and adenylate cyclase coupling (188) 243
Sandmann, J., Leissner, J., Lindmar, R. and Löffelholz, K.
The effects of phorbol esters on choline phospholipid hydrolysis in heart and brain (188) 89
Sarbin, N.S., see Gordon, J.C. (188) 313
Sauro, M.D., see Russell, D.H. (188) 139
Schalling, M., Dagerlind, Å., Goldstein, M., Ehrlich, M., Greengard, P. and Hökfelt, T.
Comparison of gene expression of the dopamine D-2 receptor and DARPP-32 in rat brain, pituitary and adrenal gland (188) 277
Scheufler, E., see Vogelgesang, R. (188) 17
Schunack, W., see Arrang, J.-M. (188) 219
Schwartz, J.-C., see Arrang, J.-M. (188) 219
Seeburg, P.H., see Lan, N.C. (188) 403
Severson, J.A., Randall, P.K. and Wilcox, R.E.
Single apomorphine pretreatment results in a rapid decline in high-affinity dopamine binding to the striatal dopamine D-2 receptor (188) 283
Shima, S.
Effects of chronic treatment with adrenocorticotropin on adenylate cyclase activity in the adrenal cortex (188) 1
Shipman, L., see Watson, S.P. (188) 273
Shreeve, S.M.
Effect of detergent solubilization on the affinity of some quinazoline derivatives for the α_1-adrenoceptor (188) 71
Skolnick, P., see Marvizón, J.-C.G. (188) 23
Sliwinski-Lis, E., see Laniyonu, A. (188) 171
Sokolovsky, M., see Galron, R. (188) 85
Sokolovsky, M., see Nadler, V. (188) 97
Soudijn, W., see Van Wijngaarden, I. (188) 301
Spierenburg, H., see Dekker, L.V. (188) 113
Su, T.-P., see McCann, D.J. (188) 211

Tang, D., see Lew, J.Y. (188) 329
Thomas, J.W., see Hood, W.F. (188) 43
Tiberi, M. and Magnan, J.
Demonstration of the heterogeneity of the κ-opioid receptors in guinea-pig cerebellum using selective and nonselective drugs (188) 379
Tiger, G., Björklund, P.-E., Cowburn, R.F., Garlind, A., O'Neill, C., Wiehager, B. and Fowler, C.J.
Effect of monovalent ions upon G proteins coupling muscarinic receptors to phosphoinositide hydrolysis in the rat cerebral cortex (188) 51
Tiong, A.H.K. and Richardson, J.S.
β-Adrenoceptor and post-receptor components show different rates of desensitization to desipramine (188) 411
Toews, M.L., see Cowlen, M.S. (188) 105
Totaro, J.A., see Pettibone, D.J. (188) 235
Tulp, M.T.M., see Van Wijngaarden, I. (188) 301

Uchida, M.-K., see Karibe, H. (188) 407
Unson, C., see Carr, C. (188) 203

Van de Velde, E., see Vanscheeuwijck, P. (188) 129
Vanscheeuwijck, P., Van de Velde, E. and Fraeyman, N.
The β-adrenergic transduction system in kidneys from young and senescent rats (188) 129
Van Wijngaarden, I., Tulp, M.T.M. and Soudijn, W.
The concept of selectivity in 5-HT receptor research (188) 301
Versteeg, D.H.G., see Dekker, L.V. (188) 113
Vogelgesang, R. and Scheufler, E.
R 56865 and flunarizine displace Ca^{2+} from phosphatidylserine monolayers in a stoichiometric manner (188) 17

Watson, S.P., Shipman, L. and Godfrey, P.P.
Lithium potentiates agonist formation of [^{3}H]CDP-diacylglycerol in human platelets (188) 273
Welk, G., see Murphy, S. (188) 399
Wiehager, B., see Tiger, G. (188) 51
Wilcox, R.E., see Severson, J.A. (188) 283
Winkler, M.E., see Yang, S.-G. (188) 289
Woyden, C.J., see Pettibone, D.J. (188) 235

Yamashita, A., Kurokawa, T., Fujii, Y., Yasuda, H. and Ishibashi, S.
Difference in sensitivity to alkaline phosphatase treatment between rat reticulocyte membranes in which β-adrenoceptor desensitization was induced by isoproterenol, dibutyryl cAMP, and phorbol ester (188) 229
Yang, S.-G., Winkler, M.E. and Hollenberg, M.D.
Contribution of the C-terminal dipeptide of transforming growth factor-α to its activity: biochemical and pharmacologic profiles (188) 289
Yara, H., Katagawa, Y., Doi, M., Ishida, T., Inoue, M., Yasuda, M., Baba, A., Iwata, H., Kitamura, K. and Sadatome, T.
An attempt to structurally convert μ-selective morphine toward δ-receptor binding: dimerization based on enkephalin conformation (188) 359
Yasuda, H., see Yamashita, A. (188) 229
Yasuda, M., see Yara, H. (188) 359

Zawadzka, H., see Lew, J.Y. (188) 329
Zorn, N.E., see Russell, D.H. (188) 139